



IMAGING AND DIAGNOSTIC TESTING

CORRELATION OF LEFT VENTRICULAR NON-COMPACTED/ COMPACTED MASS RATIO AND CLINICAL OUTCOMES IN PATIENTS WITH LEFT VENTRICULAR NON-COMPACTION

ACC Poster Contributions

Ernest N. Morial Convention Center, Hall F

Monday, April 04, 2011, 9:30 a.m.-10:45 a.m.

Session Title: MRI: Evaluation of Non-ischemic Cardiomyopathy

Abstract Category: 38. MRI

Session-Poster Board Number: 1095-183

Authors: *Yongkasem Vorasettakarnkij, Daniel J. Verdini, Peerawut Deeprasertkul, Hector M. Medina, Alexander Goehler, Elkan F. Halpern, Waleed Ahmed, Shanmugam Uthamalingam, Amparo L. Figueroa, Gyongyi P. Major, Thomas J. Brady, David E. Sosnovik, Brian B. Ghoshhajra, Godtfred Holmvang, Massachusetts General Hospital, Harvard Medical School, Boston, MA, Department of Medicine, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand*

Background: The utility of CMR in diagnosing LV non-compaction is well established. However, data is limited regarding any relationship of CMR measures of severity and cardiovascular outcomes. We hypothesize that the ratio of LV non-compacted mass (LVNCM) to compacted mass (LVCM) is correlated with major adverse cardiovascular events (MACE).

Methods: Patients referred for CMR from 01/01/2003 to 6/30/2010 who had CMR diagnosis of LV non-compaction, were included. LVEF, segmental thickness of non-compacted (NC) and compacted myocardium (C), global LVNCM and LVCM were derived. The correlations between these CMR parameters and MACE (heart failure NYHA class III-IV, thromboembolism, ventricular arrhythmia) were analyzed. Image 1 shows the non-compacted and compacted myocardium.

Results: 30 patients met entry criteria, 9 had a MACE. The LVNCM / LVCM ratio was a strong independent predictor of MACE when combined with either LVEF, total segments with NC/C > 2.3, or maximal NC/C ratio ($p = 0.008, 0.032$, and 0.028 respectively). Of these measures, the LVNCM / LVCM ratio showed the highest correlation to MACE (ROC AUC = .73, compared to .51, .63 and .64 respectively). A LVNCM/LVCM ratio > .7 predicted MACE with sensitivity, specificity, positive and negative predictive values of 89%, 43%, 40% and 90%.

Conclusions: The LVNCM/LVCM ratio shows the strongest correlation with MACE compared to LVEF, total segments with NC/C > 2.3, and maximal NC/C ratio. A ratio > .7 predicted MACE with odds ratio of 6.

